



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\therefore \frac{dV}{dx} = s - 3\pi x^2 = 0. \therefore x = \sqrt{\left(\frac{s}{3\pi}\right)}, = y.$$

Also solved by P. S. BERG, C. W. M. BLACK, J. H. DRUMMOND, A. HUME, P. H. PHILBRICK, H. C. WHITAKER, G. B. M. ZERR, and the PROPOSER.

## PROBLEMS.

20. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

$$\int_0^{\frac{1}{2}\pi} \sqrt{[(1 - e^2 \cos^2 \phi)(1 - e^2 \sin^2 \phi)]} d\phi = \text{what?}$$

21. Proposed by T. JOHN COLE, Columbus, Ohio.

In the equilateral triangle  $ABC$ ,  $AB$  the base is 10 feet. With  $B$  as a center an arc is drawn from  $C$  to  $A$ ; likewise with  $A$  as a center an arc is drawn from  $C$  to  $B$ . What is the volume of the solid generated by revolving the figure about the altitude of the triangle as an axis.

Solutions to these problems should be received on or before August 1st.

## MECHANICS.

Conducted by B. F. FINKEL, Kidder, Missouri. All Contributions to this department should be sent to him.

## SOLUTIONS TO PROBLEMS.

6. Proposed by THOMAS W. WRIGHT, M. A., Ph. D., Professor of Applied Mathematics and Physics, Union College, Schenectady, New York.

What is the effect of a charge between light and heavy cavalry, the light cavalry having the greater energy and the heavy the greater momentum?

Solution by P. H. PHILBRICK, C. E., Lake Charles, Louisiana.

Let  $M$  and  $V$  represent the mass and velocity respectively of the heavy cavalry and  $M_1$  and  $V_1$  the same of the light cavalry. Supposing the bodies to be inelastic and moving in opposite directions before impact and together after impact their common velocity after impact is,  $V = \frac{MV - M_1 V_1}{M + M_1} \dots (1)$ .

Since  $MV > M_1 V_1$  the heavy cavalry will overcome the lighter and carry it along at the above rate. The combined energy of the bodies is,  $\frac{1}{2}MV^2 + \frac{1}{2}M_1 V_1^2 \dots (2)$ . This measures the destructive effect of the charge.

7. Proposed by DE VOLSON WOOD, M. A., M. Sc., C. E., Professor of Engineering, Stevens Institute of Technology, Hoboken, New Jersey.

A hollow sphere filled with frictionless water rolls down a rough plane whose length is  $l$  and inclination  $\theta$ ; when half way down the water suddenly freezes and adheres to the sphere. Required the time of the descent.